

Geologic unit name: Disko Bugt

Geologic unit map code: PR_AE-GLMD

Unit Hierarchy:

[Disko Bugt](#)

Status: In current use

Geological period(s): Proterozoic - Eoarchaeon

Full description: The area from Disko Bugt and northwards on the Greenlandic west coast is underlain by Archean rocks that are correlated with the Rae Craton in Canada (Connelly et al., 2006; St-Onge et al., 2009). The Rae Craton of western Greenland consists of Meso- to Neoarchean orthogneiss and supracrustal complexes (see also review in Kolb et al. 2016). Dioritic orthogneiss and augen gneiss are dated at $3030 \pm 8/-5$ Ma and 2947 ± 23 Ma (U-Pb zircon), respectively. The age data are interpreted as Mesoarchean emplacement ages (Connelly et al., 2006). A felsic schist was dated at 2847 ± 4 Ma (Connelly et al., 2006). The orthogneiss consists of tonalite, trondhjemite, granodiorite, monzogranite and diorite with local leucosomes (Garde and Steenfelt, 1999). Zircons from three orthogneiss samples were dated by SHRIMP U-Pb indicating intrusion ages of ca. 2800 Ma (Nutman and Kalsbeek, 1999). An undeformed monzogranite yields 2758 ± 2 Ma as emplacement age largely postdating metamorphism and deformation (Nutman and Kalsbeek, 1999). Orthogneiss in the Rodebay area is dated at ca. 2835–2785 Ma by U-Pb zircon ages, suggesting Meso- to Neoarchean emplacement, whereas zircon rims yield ca. 2800–2760 Ma for metamorphism largely contemporaneous with granite emplacement (Connelly et al., 2006). In the Disko-Nuussuaq area, Meso- to Neoarchean orthogneiss is intercalated with greenstone belts (Garde and Steenfelt, 1999; Connelly et al., 2006). Three greenstone belts, namely Saqqaq, Itilliarsuk and Arveprinsen-Eqi greenstone belts, are distinguished (Garde and Steenfelt, 1999). The Arveprinsen-Eqi greenstone belt forms a complex regional-scale fold structure consisting of 3–4 km thick metasedimentary and metavolcanic rocks at upper greenschist to lower amphibolite facies grades. The rocks are greenstone with local pillow structures, quartz- and feldspar-phyrlic metavolcanic rock, metagabbro, BIF, phyllite and metadolerite (Garde and Steenfelt, 1999). The metagabbro forms a tholeiitic to komatiitic sill complex in the greenstone belt (Marshall and Schönwandt, 1999). Both the metagabbro and the quartz and feldspar porphyry have a volcanic arc geochemical signature (Marshall and Schönwandt, 1999; Stendal et al., 1999). The contact between orthogneiss and the Arveprinsen-Eqi and Itilliarsuk greenstone belts is intrusive, suggesting that at least these belts are Mesoarchean (Garde and Steenfelt, 1999). The Saqqaq greenstone belt forms a N 500 m thick NW-trending anticlinal structure and consists of biotite-garnet schist, amphibolite and minor ultramafic rocks (Garde et al., 1999). The Itilliarsuk greenstone belt is approx. 2.5 km thick and consists of quartz-biotite schist, staurolite-muscovite schist, metaconglomerate, amphibolite, metagabbro, BIF and calcsilicate rocks (Garde and Steenfelt, 1999; Rasmussen and Pedersen, 1999; Haugaard et al., 2013).

The Archaean basement together with Palaeoproterozoic cover rocks are variably affected by the

Palaeoproterozoic Nagssugtoqidian orogeny lasting from c. 2000 to 1750 Ma (van Gool et al. 2002). The orogen comprises dykes intruded during initial rifting, volcanic rocks formed during subduction, bodies of mantle peridotite emplaced tectonically during collision, and post-collisional granites and pegmatites. The ~2500–3600 m thick Anap Nunâ Group is situated in the northeastern Disko Bugt area and based on similar lithology correlated with the Karrat Group in the north (Garde and Steenfelt, 1999). It is weakly metamorphosed and consists of b30 m thick sandstone unconformably overlying Archean rocks at the base and in turn overlain by 50 m thick carbonate and N2000 m thick sandstone and siltstone, suggesting a shallowmarine to coastal setting progressing into a deeper basin where turbidite sequences were deposited (Garde and Steenfelt, 1999). Detrital zircons yield age groups between 3000 and 2700 Ma, and 2100 and 1900 Ma similar to the Karrat Group, confirming stratigraphic correlation (Connelly et al., 2006).

The Nuussuaq Basin covers the Disko-Svartehuk Halvø area of West Greenland. The basin is bound to the east by a system of faults, but Cretaceous-Paleogene sedimentary rocks may have extended further to the east (Chalmers et al., 1999; Pedersen et al., 2007). Approx. 6–8 km thick sedimentary units were deposited starting with Albian fluvial-lacustrine sandstone, mudstone and coal (Pedersen et al., 2007; Dam et al., 2009). The facies in the south developed into a deltaic system with sandstone and coal towards the Campanian. In the north, marine mudstone and turbidite were deposited in the deeper parts of the basin (Dam et al., 2009). Intense block faulting until the Danian divided the basin into marine turbidite-conglomerate and fluvial sandstone-mudstone sandstone-mudstone facies of largely variable thickness and distribution. Marine mudstones and lacustrine-fluvial mudstone and sandstone of upper Danian–lower Selandian are subsequently interlayered with volcanoclastic rocks marking the transition to the extensive Paleocene volcanism (Pedersen et al., 2007; Dam et al., 2009; Kolb et al. 2016).

The ca. 60 Ma volcanic rocks cover an area of ~22,000 km² on Disko, Hareø, Ubekendt Ejland, Nuussuaq and Svartehuk Halvø, and extend off-shore from 68°N to 73°N covering an area of more than 100,000 km² (Storey et al., 1998; Chalmers et al., 1999; Larsen and Pedersen, 2009). The extensive Paleocene volcanism is attributed to the impingement of the proto-Icelandic mantle plume head at the base of the lithosphere (White and McKenzie, 1989; Saunders et al., 1997). The bulk of the West Greenland volcanic rocks are of the same age as the oldest rocks of the NAIP in East Greenland, but it is unlikely that they are connected underneath the Inland Ice sheet because both units thin inland (Brooks, 2011).

The Paleocene volcanic rocks in the Nuussuaq Basin on Disko and Nuussuaq comprise the Vaigat Formation (c. 62–61 Ma) and the Maligat Formation (c. 60 Ma). The Vaigat Formation in this area is 0–1600 m thick and is dominated by olivine-rich picrites. The formation was deposited during three volcanic episodes and is divided into 10 formally defined members and about 20 informal units. The first episode gave rise to the Anaanaa Member. The second episode gave rise to the Naujanguit Member. The third episode gave rise to the Ordlingassoq Member and the minor alkaline Manitdlat Member. Contemporaneous sediments deposited during the first two episodes are the marine Eqalulik Formation, and during the third episode the nonmarine Atanikerluk Formation (Pedersen et al. 2017).

References:

- Brooks, C.K., 2011. The East Greenland rifted volcanic margin. Geological Survey of Denmark and Greenland Bulletin 24. Geological Survey of Denmark and Greenland, Copenhagen (96 pp.).
- Chalmers, J.A., Pulvertaft, T.C.R., Marcussen, C., Pedersen, A.K., 1999. New insight into the Connelly, J.N., Thrane, K., Krawiec, A.W., Garde, A.A., 2006. Linking the Palaeoproterozoic Nagsugtoqidian and Rinkian orogens through the Disko Bugt region of West Greenland. *J. Geol. Soc. Lond.* 163, 319–335.
- Dam, G., Pedersen, G.K., Sønderholm, M., Midtgaard, H.H., Larsen, L.M., Nøhr-Hansen, H., Pedersen, A.K., 2009. Lithostratigraphy of the Cretaceous-Paleocene Nuussuaq Group, Nuussuaq Basin, West Greenland. Geological Survey of Denmark and Greenland Bulletin 19. Geological Survey of Denmark and Greenland, Copenhagen, p. 171.
- Garde, A.A., Steenfelt, A., 1999. Precambrian geology of Nuussuaq and the area north-east of Disko Bugt, West Greenland. *Geology of Greenland Bulletin* 191, pp. 6–40.
- Garde, A.A., Thomassen, B., Tukiainen, T., Steenfelt, A., 1999. A gold-bearing volcanogenic-exhalative horizon in the Archaean(?) Saqqaq supracrustal rocks, Nuussuaq, West Greenland. *Geology of Greenland Bulletin* 181, pp. 119–128.
- Haugaard, R., Frei, R., Stendal, H., Konhauser, K., 2013. Petrology and geochemistry of the ~2.9 Ga Itilliarsuk banded iron formation and associated supracrustal rocks, West Greenland: source characteristics and depositional environment. *Precambrian Res.* 229, 150–176.
- Kolb, J., Keiding, J.K., Steenfel, A., Secher, K., Keulen, N., Rosa N.D., Stensgaard, B.M. 2016. Metallogeny of Greenland. *Ore Geology Reviews* 78, 493–555.
- Marshall, B., Schønswandt, H.K., 1999. An Archaean sill complex and associated supracrustal rocks, Arveprinsens Ejland, north-east Disko Bugt, West Greenland. *Geology of Greenland Survey Bulletin* 181, pp. 87–102.
- Nutman, A.P., Kalsbeek, F., 1999. SHRIMP U-Pb zircon ages for Archaean granitoid rocks, Pedersen, A. K., Larsen, L. M., & Pedersen, G. K. (2017). Lithostratigraphy, geology and geochemistry of the volcanic rocks of the Vaigat Formation on Disko and Nuussuaq, Paleocene of West Greenland. Copenhagen: The Geological Survey of Denmark and Greenland. Geological Survey of Denmark and Greenland (GEUS) Bulletin, Vol.. 39: 251 pp.
- Pedersen, A.K., Pedersen, G.K., Larsen, L.M., Pulvertaft, T.C.R., Sønderholm, M., Dueholm, K.S., 2007. Geological Map of the South-East Coast of Nuussuaq Between Ataata Kuua and Saqqaq dalen, Central West Greenland, 1:50 000, With Detailed Sections. Geological Survey of Denmark and Greenland, Copenhagen.
- Rasmussen, H., Pedersen, L.F., 1999. Stratigraphy, structure and geochemistry of Archaean supracrustal rocks from Oqaatsut and Naajaat Qaqqaat, north-east Disko Bugt, West Greenland. *Geology of Greenland Bulletin* 181, pp. 65–78.
- Saunders, A.D., Fitton, J.G., Kerr, A.C., Norry, M.J., Kent, R.W., 1997. The North Atlantic Igneous province. In: Mahoney, J.J., Coffin, M.F. (Eds.), *Large Igneous Provinces*. American Geophysical Union Geophysical Monograph 100, pp. 45–93 (Washington D.C.).
- St-Onge, M.R., van Gool, J.A.M., Garde, A.A., Scott, D.J., 2009. Correlation of Archaean and Palaeoproterozoic units between northeastern Canada and western Greenland: constraining the pre-

- collisional upper plate accretionary history of the Trans-Hudson orogen. In: Cawood, P.A., Kröner, A. (Eds.), *Earth Accretionary Systems in Space and Time*. The Geological Society of London, Special Publications 318, pp. 193–235 (London).
- Stendal, H., Knudsen, C., Marker, M., Thomassen, B., 1999. Gold mineralisation at Eqi, north-east Disko Bugt, West Greenland. *Geology of Greenland Bulletin* 181, pp. 129–140.
- Storey, M., Duncan, R.A., Pedersen, A.K., Larsen, L.M., Larsen, H.C., 1998. Ar/Ar geochronology of the West Greenland Tertiary volcanic province. *Earth Planet. Sci. Lett.* 160, 569–586.
- White, D., McKenzie, D., 1989. Magmatism at rift zones: the generation of volcanic continental margins and flood basalts. *J. Geophys. Res. Solid Earth* 94, 7685–7729.
- van Gool, J.A.M., Connelly, J.N., Marker, M. & Mengel, F.[C.] 2002: The Nagssugtoqidian Orogen of West Greenland: tectonic evolution and regional correlations from a West Greenland perspective. *Canadian Journal of Earth Sciences* 39, 665–686.